arXiv:1502.02188v3 [cs.IT] 9 Mar 2015

Entropy-power inequality for weighted entropy

Y. Suhov$^{1-3}$, S. Yasaei Sekeh^{4}, M. Kelbert^{5,6}

[1] DPMMS, University of Cambridge, UK
[2] Math Dept, Penn State University, PA, USA
[3] IPIT RAS, Moscow, RF
[4] DEs, Federal University of São Carlos, SP, Brazil
[5] Dept of Mathematics, University of Swansea, UK
[6] Moscow Higher School of Economics, RF

Abstract

We analyse an analog of the entropy-power inequality for the weighted entropy. In particular, we discuss connections with weighted Lieb‘s splitting inequality and an Gaussian additive noise formula. Examples and counterexamples are given, for some classes of probability distributions.

Key words: weight function, weighted differential entropy, weighted entropy-power inequality, weighted Lieb′s splitting inequality, weighted conditional and mutual entropies, additive Gaussian noise,

2000 MSC: 60A10, 60B05, 60C05

1 Introduction. The weighted entropy-power inequality

Let $x \in \mathbb{R} \mapsto \phi(x) \geq 0$ be a given (measurable) function. The weighted differential entropy (WDE) $h^{\rm w}_\phi(Z)$ of a real-valued random variable (RV) Z with a probability density function (PDF) f_Z is defined by the formula

$$h^{\rm w}_\phi(Z) = h^{\rm w}_\phi(f_Z) := -\mathbb{E}\,\phi(Z)\ln f_Z(Z) = -\int_{\mathbb{R}^n} \phi(x) f_Z(x) \ln f_Z(x) {\rm d}x, \tag{1.1}$$

assuming that the integral is absolutely convergent (with the usual agreement that $0 \cdot \ln 0 = 0$). Cf. [1], [2], [7]. For $\phi(x) \equiv 1$, the definition yields the standard (Shannon) differential entropy (SDE). Furthermore, ϕ is called a weight function (WF). When we say that $h^{\rm w}_\phi(Z)$ is finite we mean that RV Z has a PDF f_Z, and the integral in (1.1) absolutely converges.

We propose the following bound which we call the weighted entropy-power inequality (WEPI): for two independent RVs X_1 and X_2, with $X = X_1 + X_2$,

$$\exp\left[\frac{2\,h^{\rm w}_\phi(X_1)}{\mathbb{E}\,\phi(X_1)}\right] + \exp\left[\frac{2\,h^{\rm w}_\phi(X_2)}{\mathbb{E}\,\phi(X_2)}\right] \leq \exp\left[\frac{2\,h^{\rm w}_\phi(X)}{\mathbb{E}\,\phi(X)}\right], \tag{1.2}$$

assuming that the WDEs $h^{\rm w}_\phi(X)$, $h^{\rm w}_\phi(X_1)$ and $h^{\rm w}_\phi(X_2)$ are finite, as well as the expected values $\mathbb{E}\,\phi(X), \mathbb{E}\,\phi(X_1), \mathbb{E}\,\phi(X_2)$ (the latter means that $\mathbb{E}\,\phi(X), \mathbb{E}\,\phi(X_1), \mathbb{E}\,\phi(X_2) \in (0,\infty)$). Again, for $\phi(x) \equiv 1$, this yields the famous EPI put forward by Shannon; see [3], [8], [4], [5]. In this note we offer a sufficient condition for (1.2) (see Eqns (1.5) and (1.6) below); the origins of bound (1.6) go back to Ref. [6]. We set:

$$\alpha = \tan^{-1}\left(\exp\left[\frac{h^{\rm w}_\phi(X_2)}{\mathbb{E}\,\phi(X_2)} - \frac{h^{\rm w}_\phi(X_1)}{\mathbb{E}\,\phi(X_1)}\right]\right),\ Y_1 = \frac{X_1}{\cos\alpha},\ Y_2 = \frac{X_2}{\sin\alpha}, \tag{1.3}$$

and

$$\kappa = \exp\left[\frac{2h^{\rm w}_\phi(X_1)}{\mathbb{E}\,\phi(X_1)}\right] + \exp\left[\frac{2h^{\rm w}_\phi(X_2)}{\mathbb{E}\,\phi(X_2)}\right]. \tag{1.4}$$

Theorem 1: *Given independent RVs X_1, X_2 and a WF ϕ, set $X = X_1 + X_2$ and make the following suppositions.* (i) *The expected values obey*

$$\begin{array}{l}\mathbb{E}\,\phi(X_1) \geq \mathbb{E}\,\phi(X) \ \text{ and } \ \mathbb{E}\,\phi(X_2) \geq \mathbb{E}\,\phi(X) \ \text{ if } \kappa \geq 1,\\ \mathbb{E}\,\phi(X_1) \leq \mathbb{E}\,\phi(X) \ \text{ and } \ \mathbb{E}\,\phi(X_2) \leq \mathbb{E}\,\phi(X) \ \text{ if } \kappa \leq 1.\end{array} \tag{1.5}$$

(ii) *With α and Y_2, Y_2 as in Eqn* (1.3),

$$(\cos\alpha)^2\ h^{\rm w}_{\phi_{\rm C}}(Y_1) + (\sin\alpha)^2\ h^{\rm w}_{\phi_{\rm S}}(Y_2) \leq h^{\rm w}_\phi(X). \tag{1.6}$$

Here

$$\phi_{\rm C}(x) = \phi(x\cos\alpha), \quad \phi_{\rm S}(x) = \phi(x\sin\alpha) \tag{1.7}$$

and we assume finite WDEs $h^{\rm w}_\phi(X)$ and

$$h^{\rm w}_{\phi_{\rm C}}(Y_2) = -\mathbb{E}\phi_{\rm C}(Y_2)\ln f_{Y_2}(Y_2), \quad h^{\rm w}_{\phi_{\rm S}}(Y_2) = -\mathbb{E}\phi_{\rm S}(Y_2)\ln f_{Y_2}(Y_2).$$

Then WEPI (1.2) *holds true.*

Proof: We can write

$$h^{\rm w}_\phi(X_1) = h^{\rm w}_{\phi_{\rm C}}(Y_2) + \mathbb{E}\,\phi(X_1)\log\ \cos\alpha, \quad h^{\rm w}_\phi(X_2) = h^{\rm w}_{\phi_{\rm S}}(Y_2) + \mathbb{E}\,\phi(X_2)\log\ \sin\alpha.$$

Using (1.6), we have the following inequality:

$$\begin{aligned} h^{\rm w}_\phi(X) \geq (\cos\alpha)^2\Big[h^{\rm w}_\phi(X_1) - \mathbb{E}\,\phi(X_1)\log\ \cos\alpha\Big]\\ +(\sin\alpha)^2\Big[h^{\rm w}_\phi(X_2) - \mathbb{E}\,\phi(X_2)\log\ \sin\alpha\Big].\end{aligned}$$

Furthermore, recalling (1.4) we obtain:

$$\begin{aligned} & h^{\mathrm{w}}_\phi(X) \\ & \geq \frac{1}{2\kappa}\Big[\mathbb{E}\phi(X_1)\log\,\kappa\Big]\exp\left[\frac{2h^{\mathrm{w}}_\phi(X_1)}{\mathbb{E}\,\phi(X_1)}\right]+\frac{1}{2\kappa}\Big[\mathbb{E}\,\phi(X_2)\log\,\kappa\Big]\exp\left[\frac{2h^{\mathrm{w}}_\phi(X_2)}{\mathbb{E}\,\phi(X_2)}\right]. \end{aligned}$$

By virtue of assumption (1.5), we derive:

$$h^{\mathrm{w}}_\phi(X)\geq\frac{1}{2}\mathbb{E}\,\phi(X)\log\,\kappa.$$

The definition of κ in Eqn (1.4) leads directly to the result. $\square$

Paying homage to Ref. [6], we call the bound (1.6) the WLSI (weighted Lieb′s splitting inequality). In the spirit of [6], the following Theorem 2 can be offered. (The notation used in Theorem 2 is self-explanatory; the proof of Theorem 2 is one-line and omitted.)

Theorem 2: *Let f and g be PDFs on $\mathbb{R}$ and ϕ a given WF. Assume that the WDEs $h^{\mathrm{w}}_\phi(f*g)$, $h^{\mathrm{w}}_\phi(f)$ and $h^{\mathrm{w}}_\phi(g)$ are finite, as well as expected values $\mathbb{E}_f\,\phi$, $\mathbb{E}_g\,\phi$. Set*

$$\tau=\exp\left[\frac{2h^{\mathrm{w}}_\phi(f)}{\mathbb{E}_f\,\phi}\right]+\exp\left[\frac{2h^{\mathrm{w}}_\phi(g)}{\mathbb{E}_g\,\phi}\right].$$

Also suppose that

$$\begin{aligned} &\mathbb{E}_f\,\phi\geq\mathbb{E}_{f*g}\,\phi\ \text{ and }\ \mathbb{E}_g\,\phi\geq\mathbb{E}_{f*g}\,\phi\ \text{ if }\tau\geq 1,\\ &\mathbb{E}_f\,\phi\leq\mathbb{E}_{f*g}\,\phi\ \text{ and }\ \mathbb{E}_g\,\phi\leq\mathbb{E}_{f*g}\,\phi\ \text{ if }\tau\leq 1. \end{aligned}\tag{1.8}$$

and the following inequality holds:

$$\begin{aligned} 2h^{\mathrm{w}}_\phi(f*g)\geq{}&2\lambda h^{\mathrm{w}}_\phi(f)+2(1-\lambda)h^{\mathrm{w}}_\phi(g)\\ &-\mathbb{E}_f\,\phi\lambda\log\lambda-\mathbb{E}_g\,\phi(1-\lambda)\log(1-\lambda), \end{aligned}\tag{1.9}$$

where $\lambda\in[0,1]$ is given by

$$\lambda=\tau^{-1}\,\exp\left[\frac{2h^{\mathrm{w}}_\phi(f)}{\mathbb{E}_f\,\phi}\right].\tag{1.10}$$

Then Eqn (1.2) holds true for independent RVs X_1 and X_2 where $X_1\sim f$, $X_2\sim g$.

Remark. The arguments developed in Section 1 do not use the fact that RVs X_1 and X_2 possess PDFs. The question of whether the WEPI (as it is presented in Eqn (1.2) or in a modified form) may hold for cases of discrete distributions requires a separate investigation. However, constructions used in Section 3 demand existence of PDFs f_{X_1} and f_{X_2} although some of their technical parts are valid in a more general situation.

2 Examples and counterexamples

In this section we give several examples where the above inequalities hold or do not hold true.

2.1. Examples. First, let us discuss specific conditions equivalent to (1.2), (1.5) or (1.6), for various pairs of RVs. In the next subsection we present results of numerical simulations showing domains of parameters where Eqns (1.2), (1.5) and (1.6) are fulfilled or violated.

2.1.1. (Normal distributions) Let X_1, X_2 be two independent normal RVs: $X_1 \sim \mathrm{N}(0,\sigma_1^2)$, $X_2 \sim \mathrm{N}(0,\sigma_2^2)$ and $X \sim \mathrm{N}(0,\sigma_1^2+\sigma_2^2)$. Recall (see [7], Example 3.1), the WDE $h^{\mathrm{w}}_{\phi}(Z)$ of a normal random variable $Z \sim \mathrm{N}(0,\sigma^2)$ reads

$$h^{\mathrm{w}}_{\phi}(Z) = \frac{\log\left(2\pi\sigma^2\right)}{2}\mathbb{E}\phi(Z) + \frac{\log e}{2\sigma^2}\mathbb{E}Z^2\phi(Z);$$

we will use it for $Z = X, X_1, X_2$. The condition $\kappa \geq (\leq) 1$ is re-written as

$$\sigma_1^2 \exp\left\{\frac{\mathbb{E}[X_1^2\phi(X_1)]}{\sigma_1^2\,\mathbb{E}\,\phi(X_1)}\right\} + \sigma_2^2 \exp\left\{\frac{\mathbb{E}[X_2^2\phi(X_2)]}{\sigma_2^2\,\mathbb{E}\,\phi(X_2)}\right\} \geq (\leq)\,(2\pi)^{-1}. \tag{2.1}$$

We have to match it with inequalities

$$\mathbb{E}\phi(X_1),\ \ \mathbb{E}\phi(X_2) \geq (\leq)\mathbb{E}\,\phi(X)$$

to fulfill (1.5)

To specify the WLSI (1.6), we write:

$$h^{\mathrm{w}}_{\phi_{\mathrm{C}}}(Y_1) = \frac{1}{2}\left[\log\frac{2\pi\sigma_1^2}{(\cos\alpha)^2}\right]\mathbb{E}\,\phi_{\mathrm{C}}(Y_1) + \frac{(\cos\alpha)^2\log e}{2\sigma_1^2}\,\mathbb{E}\big[Y_1^2\phi_{\mathrm{C}}(Y_1)\big]. \tag{2.2}$$

Pluging-in the definition of ϕ_{C}:

$$\mathbb{E}\phi_{\mathrm{C}}(Y_1) = \mathbb{E}\phi(X_1), \quad \mathbb{E}\big[Y_1^2\phi_{\mathrm{C}}(Y_1)\big] = \frac{\mathbb{E}\big[X_1^2\phi(X_1)\big]}{(\cos\alpha)^2}.$$

Similar equations hold for $h^{\mathrm{w}}_{\phi_{\mathrm{S}}}(Y_2)$. Then Eqn (1.6) takes the form

$$\begin{aligned}&\Big[\log\Big(2\pi(\sigma_1^2+\sigma_2^2)\Big)\Big]\mathbb{E}\phi(X) + \frac{\log e}{\sigma_1^2+\sigma_2^2}\,\mathbb{E}\big[X^2\phi(X)\big]\\ &\geq (\cos\alpha)^2\left[\log\frac{2\pi\sigma_1^2}{(\cos\alpha)^2}\right]\mathbb{E}\phi(X_1) + \frac{(\cos\alpha)^2\log e}{\sigma_1^2}\,\mathbb{E}[X_1^2\phi(X_1)]\\ &\quad+(\sin\alpha)^2\left[\log\frac{2\pi\sigma_2^2}{(\sin\alpha)^2}\right]\mathbb{E}\phi(X_2) + \frac{(\sin\alpha)^2\log e}{\sigma_2^2}\,\mathbb{E}[X_2^2\phi(X_2)].\end{aligned} \tag{2.3}$$

2.1.2. (Gamma-distributions) Let X_1 and X_2 have Gamma distributions, with PDFs $f_{X_i}(x) = \dfrac{\lambda^{\beta_i}}{\Gamma(\beta_i)}x^{\beta_i-1}e^{-\lambda x}$, $i=1,2$, and $f_X(x) = \dfrac{\lambda^{\beta}}{\Gamma(\beta)}x^{\beta-1}e^{-\lambda x}$, $x>0$ where $\beta = \beta_1+\beta_2$. The WDEs are

$$h^{\mathrm{w}}_{\phi}(X_i) = (1-\beta_i)\mathbb{E}[\phi(X_i)\log X_i] + \lambda\mathbb{E}[X_i\phi(X_i)] + \log\left(\frac{\Gamma(\beta_i)}{\lambda^{\beta_i}}\right)\mathbb{E}\phi(X_i)$$

and similarly for X (with β instead of β_i). The condition $\kappa \geq (\leq) 1$ reads

$$\begin{aligned}&\left(\frac{\Gamma\beta_1}{\lambda^{\beta_1}}\right)\exp\left\{\frac{\lambda\mathbb{E}[X_1\phi(X_1)]-(\beta_1-1)\mathbb{E}[\phi(X_1)\log X_1]}{\mathbb{E}\,\phi(X_1)}\right\}\\&\quad+\left(\frac{\Gamma\beta_2}{\lambda^{\beta_2}}\right)\exp\left\{\frac{\lambda\mathbb{E}[X_2\phi(X_2)]-(\beta_2-1)\mathbb{E}[\phi(X_2)\log X_2]}{\mathbb{E}\,\phi(X_2)}\right\}\geq(\leq)\,1;\end{aligned}\tag{2.4}$$

as above, it has to be in conjunction with $\mathbb{E}\,\phi(X_1), \mathbb{E}\,\phi(X_2) \geq (\leq)\, \mathbb{E}\,\phi(X)$. The WLSI (1.6) takes the following form:

$$\begin{aligned}&\log\frac{\Gamma(\beta)}{\lambda^{\beta}}\mathbb{E}\,\phi(X)-(\beta-1)\mathbb{E}[\phi(X)\log X]+\lambda\mathbb{E}[X\phi(X)]\\&\quad\geq\big(\cos\alpha\big)^2\Big[\lambda\mathbb{E}[X_1\phi(X_1)]-(\beta_1-1)\mathbb{E}[\phi(X_1)\log X_1]\Big]+\mathbb{E}\,\phi(X_1)\log\frac{\Gamma(\beta_1)}{\lambda^{\beta_1}\cos\alpha}\\&\qquad+\big(\sin\alpha\big)^2\Big[\lambda\mathbb{E}[X_2\phi(X_2)]-(\beta_2-1)\mathbb{E}[\phi(X_2)\log X_2]\Big]+\mathbb{E}\,\phi(X_2)\log\frac{\Gamma(\beta_2)}{\lambda^{\beta_2}\sin\alpha}.\end{aligned}\tag{2.5}$$

2.1.3. (Exponential distributions) Let X_1 and X_2 be two independent exponential RVs, with means λ_1^{-1} and λ_2^{-1}, and WDEs $h^{\rm w}_{\phi}(X_i) = \big(\lambda_i \log\, \lambda_i\big)\mathbb{E}\phi(X_i) + \mathbb{E}X_i\phi(X_i)$, $i = 1, 2$. See [7], Example 3.2. Then $f_X(x) = \dfrac{\lambda_1\lambda_2}{\lambda_1-\lambda_2}\big(e^{-\lambda_2 x} - e^{-\lambda_1 x}\big)$. The inequality $\kappa \geq (\leq) 1$ becomes

$$\lambda_2^2\exp\left\{\frac{2\lambda_1\mathbb{E}[X_1\phi(X_1)]}{\mathbb{E}\,\phi(X_1)}\right\}+\lambda_1^2\exp\left\{\frac{2\lambda_2\,\mathbb{E}_{X_2}[X_2\phi]}{\mathbb{E}\,\phi(X_2)}\right\}\geq(\leq)\,\lambda_1^2\lambda_2^2,\tag{2.6}$$

and to fulfill Eqn (1.5) it has to be combined with

$$\mathbb{E}\,\phi(X_1),\ \mathbb{E}\,\phi(X_2)\geq(\leq)\,\frac{\lambda_1\mathbb{E}\,\phi(X_2)-\lambda_2\mathbb{E}\,\phi(X_1)}{\lambda_1-\lambda_2}.\tag{2.7}$$

In this example, the WLSI (1.6) reads

$$\begin{aligned}&\frac{\log\ \lambda_1\lambda_2}{\lambda_1-\lambda_2}\Big\{\lambda_2\mathbb{E}\phi(X_1)-\lambda_1\mathbb{E}\phi(X_2)\Big\}+\frac{\lambda_2}{\lambda_1-\lambda_2}\mathbb{E}\left[\phi(X_1)\log\left(\frac{e^{-\lambda_2X_1}-e^{-\lambda_1X_1}}{\lambda_1-\lambda_2}\right)\right]\\&\qquad-\frac{\lambda_1}{\lambda_1-\lambda_2}\mathbb{E}\left[\phi(X_2)\log\ \left(\frac{e^{-\lambda_2X_2}-e^{-\lambda_1X_2}}{\lambda_1-\lambda_2}\right)\right]\\&\quad\geq\lambda_1(\cos\alpha)^2\,\mathbb{E}[X_1\phi(X_1)]-(\cos\alpha)^2\mathbb{E}\,\phi(X_1)\log(\lambda_1\cos\alpha)\\&\qquad\qquad+\lambda_2(\sin\alpha)^2\mathbb{E}[X_2\phi(X_2)]-(\sin\alpha)^2\mathbb{E}\,\phi(X_2)\log(\lambda_2\sin\alpha).\end{aligned}\tag{2.8}$$

2.1.4. (Uniform distributions) Set $\Phi(x) = \int\limits_0^x \phi(u){\rm d}u$, $\Phi^*(x) = \int\limits_0^x u\,\phi(u){\rm d}u$. Let $X_1 \sim {\rm U}(a_1, b_1)$ and $X_2 \sim {\rm U}(a_2, b_2)$, independently, where $L_i := b_i - a_i > 0$, $i = 1, 2$. The WDE $h^{\rm w}_{\phi}(X_i) = \dfrac{\Phi(b_i)-\Phi(a_i)}{L_i}\log\, L_i$. Suppose for definiteness that $L_2 \geq L_1$ or, equivalently, $C_1 := a_2 + b_1 \leq$

$a_1 + b_2 =: C_2$. Then PDF f_X for $X = X_1 + X_2$ has a trapezoidal form with corner points at $A = a_1 + a_2$, C_1, C_2 and $B = b_1 + b_2$:

$$f_X(x) = \frac{1}{L_1 L_2} \times \begin{cases} 0, & \text{if } x < A \text{ or } x > B, \\ x - A, & \text{if } A < x < C_1, \\ L_1, & \text{if } C_1 < x < C_2, \\ B - x, & \text{if } C_2 < x < B. \end{cases}$$

The condition $\kappa \geq (\leq) 1$ takes the form

$$L_1^2 + L_2^2 \geq (\leq)\, 1. \tag{2.9}$$

Consider the quantity Λ (which may be as positive as well as negative):

$$\begin{aligned} \Lambda = {} & \frac{\log L_1}{L_2}\Big[\Phi(C_1) - \Phi(C_2)\Big] \\ & + \frac{1}{L_1 L_2}\left[\int_A^{C_1} \phi(x)(x-A)\, \log(x-A)\,\mathrm{d}x + \int_{C_2}^{B} \phi(x)(B-x)\, \log(B-x)\mathrm{d}x\right]. \end{aligned} \tag{2.10}$$

Then

$$\begin{aligned} \mathbb{E}\,\phi(X) = {} & \frac{1}{L_1 L_2}\Bigg\{\Big[\Phi^*(C_1) - \Phi^*(A) - \Phi^*(B) + \Phi^*(C_2)\Big] \\ & -A\Big[\Phi(C_1) - \Phi(A)\Big] + L_1\Big[\Phi(C_2) - \Phi(C_1)\Big] + B\Big[\Phi(B) - \Phi(C_2)\Big]\Bigg\}. \end{aligned}$$

To satisfy condition (1.5), we have to assume that

$$L_2\big[\Phi(b_1) - \Phi(a_1)\big],\ L_1\big[\Phi(b_2) - \Phi(a_2)\big] \geq (\leq)\, \mathbb{E}\,\phi(X), \tag{2.11}$$

in conjunction with bound (2.9).

The WLSI (1.6) becomes

$$\begin{aligned} h^{\mathrm{w}}_\phi(X) &= -\Lambda + \big[\log\big(L_1 L_2\big)\big]\mathbb{E}\,\phi(X) \\ &\geq (\cos\alpha)^2 \frac{\Phi(b_1) - \Phi(a_1)}{L_1} \log\frac{L_1}{\cos\alpha} + (\sin\alpha)^2 \frac{\Phi(b_2) - \Phi(a_2)}{L_2} \log\frac{L_2}{\sin\alpha}. \end{aligned} \tag{2.12}$$

2.1.5. (A mixed case) Let X_1 be a Gamma RV with PDF $f_{X_1}(x) = \dfrac{\lambda^\beta}{\Gamma(\beta)} x^{\beta-1} e^{-\lambda x}$ and the cumulative distribution function $F_{X_1}(x)$. The WDE $h^{\mathrm{w}}_\phi(X_1)$ has been specified in Example 2.1.3. Take RV X_2 from the uniform distribution $\mathrm{U}(a,b)$, where $L := b - a > 0$, independent of X_1. The WDE $h^{\mathrm{w}}_\phi(X_1)$ has been specified in Example 2.1.4. We can write

$$f_X(t) = \frac{1}{L}\Big[F_{X_1}(t-a) - F_{X_1}(t-b)\Big].$$

As in Example 2.1.3, let $\Phi(x)=\int_0^x\phi(u)\mathrm{d}u$. Next, set:

$$\Theta=\int_0^\infty\phi(x)\left[F_{X_1}(x-a)-F_{X_1}(x-b)\right]\log\Big[F_{X_1}(x-a)-F_{X_1}(x-b)\Big]\,\mathrm{d}x. \tag{2.13}$$

Then

$$\begin{aligned}h^{\rm w}_\phi(X)=&\frac{\log(L\,\Gamma(\beta))}{L}\\&\times\mathbb{E}\Big[\Phi(X_1+b)\mathbf{1}(X_1>-b)-\Phi(X_1+a)\mathbf{1}(X_1>-a)\Big]-\frac{\Theta}{L}.\end{aligned} \tag{2.14}$$

The quantity κ is specified by

$$\kappa=\left(\frac{\Gamma(\beta)}{\lambda^\beta}\right)^2\exp\Big[\frac{\lambda\mathbb{E}[X_1\phi(X_1)]-(\beta-1)\mathbb{E}[\phi(X_1)\log X_1]}{\mathbb{E}\,\phi(X_1)}\Big]+L^2. \tag{2.15}$$

Note that if $b-a\geq 1$ we always have $\kappa>1$. To fulfill condition (1.5), we have to assume that

$$\begin{cases}L\mathbb{E}\,\phi(X_1)\geq(\leq)\,\mathbb{E}[\Phi(X_1+b)-\Phi(X_1+a)]\\ \Phi(b)-\Phi(a)\geq(\leq)\,\mathbb{E}[\Phi(X_1+b)-\Phi(X_1+a)]\end{cases}\text{depending on }\kappa\geq(\leq)1. \tag{2.16}$$

The WLSI (1.6) takes the form:

$$\begin{aligned}h^{\rm w}_\phi(X):=&\log[L\Gamma(\beta)]\,\mathbb{E}\Big[\Phi(X_1+b)\mathbf{1}(X_1>-b)-\Phi(X_1+a)\mathbf{1}(X_1>-a)\Big]-\Theta\\ \geq&L(\cos\alpha)^2\left[(1-\beta)\mathbb{E}[\phi(X_1)\log X_1]+\lambda\mathbb{E}[X_1\phi(X_1)]+\log\left(\frac{\Gamma(\beta)}{\lambda^\beta\cos\alpha}\right)\mathbb{E}\phi(X_1)\right]\\&+(\sin\alpha)^2\frac{\Phi(b)-\Phi(a)}{b-a}\log\left(\frac{L}{\sin\alpha}\right).\end{aligned} \tag{2.17}$$

2.1.6. (Cauchy distributions) Let X_1, X_2 be independent, with PDFs $f_{X_j}(x)=(\pi\theta_j)^{-1}\times\Big[1+(x-\mu_j)^2/\theta_j^2\Big]^{-1}$, $x\in\mathbb{R}$, $j=1,2$. Then $f_{X_j}(x)$ is f the same form, with parameters $\mu=\mu_1+\mu_2$ and $\theta=\theta_1+\theta_2$. For the WDEs $h^{\rm w}_\phi(X)$ we have the formula

$$h^{\rm w}_\phi(X)=\mathbb{E}\,\phi(X)\log(\pi\theta)+\mathbb{E}\left[\phi(X)\log(1+(X-\mu)^2/\theta^2)\right]$$

and similarly for $h^{\rm w}_\phi(X_i)$. The condition $\kappa\geq(\leq)1$ is re-written as

$$\begin{aligned}&\theta_1^2\exp\Big\{\frac{2\mathbb{E}[\phi(X_1)\log(1+(X_1-\mu_1)^2/\theta_1^2)]}{\mathbb{E}\,\phi(X_1)}\Big\}\\&\quad+\theta_2^2\,\exp\Big\{\frac{2\mathbb{E}[\phi(X_2)\log(1+(X_2-\mu_2)^2/\theta_2^2)]}{\mathbb{E}\,\phi(X_2)}\Big\}\geq(\leq)\,\pi^{-2};\end{aligned} \tag{2.18}$$

to satisfy (1.6) we have to match it with $\mathbb{E}\phi(X_1)$, $\mathbb{E}\phi(X_2)\geq(\leq)\mathbb{E}\,\phi(X)$. The WLSI (1.6) reads

$$\begin{aligned}&\mathbb{E}\,\phi(X)\log(\pi\theta)+\mathbb{E}\left[\phi(X)\log(1+(X-\mu)^2/\theta^2)\right]\\&\quad\geq(\cos\alpha)^2\mathbb{E}\left[\phi(X_1)\log(1+(X_1-\mu_1)^2/\theta_1^2\right]+(\cos\alpha)^2\mathbb{E}\,\phi(X_1)\log\frac{\pi\theta_1}{\cos\alpha}\\&\qquad+(\sin\alpha)^2\mathbb{E}\left[\phi(X_2)\log(1+(X_2-\mu_2)^2/\theta_2^2)\right]+(\sin\alpha^2)\mathbb{E}\,\phi(X_2)\log\frac{\pi\theta_2}{\sin\alpha}.\end{aligned} \tag{2.19}$$

2.1.7. (A distribution with an infinite SDE) Here we take independent $X_1, X_2 \sim g$ where g is a PDF on $(1,\infty)$:

$$g(x)=\frac{1}{x\big[(\ln x)^2+1\big]},\quad x>1.$$

Here we have that

$$f_X(t)=\int g(t-s)g(s)\mathrm{d}s=\int_1^{t-1}\frac{1}{(t-s)\Big[1+(\ln(t-s))^2\Big]s\Big[1+(\ln s)^2\Big]}\mathrm{d}s.$$

Therefore

$$\begin{aligned}&h^{\mathrm{w}}_{\phi}(X)\\&=-\int_2^{\infty}\phi(x)\int_1^{x-1}\frac{1}{(x-s)\Big[1+(\ln(x-s))^2\Big]s\Big[1+(\ln s)^2\Big]}\mathrm{d}s\\&\qquad\times\log\left(\int_1^{x-1}\frac{1}{(x-w)\Big[1+(\ln(x-w))^2\Big]w\Big[1+(\ln w)^2\Big]}\mathrm{d}w\right)\mathrm{d}x,\end{aligned}\tag{2.20}$$

assuming that WF ϕ decreases fast enough so that the integral in (2.20) absoltely converges. The bound $\kappa\geq(\leq)\,1$ now reads:

$$\exp\left\{\frac{2\mathbb{E}\Big[\phi(X_1)\log\big\{X_1[1+(\ln X_1)^2+1]\big\}\Big]}{\mathbb{E}\,\phi(X_1)}\right\}\geq(\leq)\,\frac{1}{2}\,,\tag{2.21}$$

and we again have to match it with inequality $\mathbb{E}\phi(X_1),\ \mathbb{E}\phi(X_2)\geq(\leq)\mathbb{E}\,\phi(X)$.

The WLSI (1.6) takes the form:

$$\begin{aligned}&-\int_2^{\infty}\phi(x)\int_1^{x-1}\frac{1}{(x-s)\Big[1+(\ln(x-s))^2\Big]s\Big[1+(\ln s)^2\Big]}\mathrm{d}s\\&\qquad\times\log\left(\int_1^{x-1}\frac{1}{(x-w)\Big[1+(\ln(x-w))^2\Big]w\Big[1+(\ln w)^2\Big]}\mathrm{d}w\right)\mathrm{d}x\\&\qquad\geq\mathbb{E}\big[\phi(X_1)\log X_1\big]+\mathbb{E}\Big\{\phi(X_1)\log\big[1+(\ln X_1)^2)\big]\Big\}\\&\qquad\qquad-\mathbb{E}\,\phi(X_1)\Big[(\cos\alpha)^2\log\cos\alpha+(\sin\alpha)^2\log\sin\alpha\Big].\end{aligned}\tag{2.22}$$

2.2. Numerical results. As was said, in this subsection we comment on some numerical evidence that (i) Eqn (1.2) does not always hold true, and (ii) assumptions (1.5) (1.6) in Theorem 1 are not necessary for the WEPI (1.2). Our observations are of a preliminary character, and we think that further numerical simulations are needed here, to build a detailed picture.

2.2.1. (Normal distributions) Assume that X_1 and X_2 are normal RVs as in Example 2.1.1. Choose $\phi(x) = |x^2 - 2|$. The graph in Figure 2.2.1 presetns the difference between the RHS and the LHS in (1.2): when this difference is non-negative, the WEPI is satisfied, otherwise the WEPI fails.The graph shows that there is a domain of parameter (σ_1, σ_2) where Eqn (1.2) does not hold. Additional simulations state that there is a domain where (1.2) holds and only one of Eqns (1.5), (1.6) is satisfied. For example, in the square where $0.55 < \sigma_1, \sigma_2 < 0.551$ the condition (1.6) is violated but (1.5), (1.2) hold true.

2.2.2. (Gamma distributions) Assume that X_1 and X_2 are gamma RVs as in Example 2.1.2. Here we choose $\phi(x) = xe^{-x}$. The graph in Figure 2.2.2 again shows the difference between the RHS and the LHS in (1.2): Here the WEPI is satisfied (in the presented range of parameters (β_1, β_2)). As in Example 2.2.1, additional simulations assert that there is a domain where (1.2) holds while none of (1.5), (1.6) is satisfied. For example, in the square $0.01 < \beta_1 < 0.1$ and $5 < \beta_2 < 6.1$ both conditions (1.5), (1.6) are violated but (1.2) holds true.



Figure 2.2.1



Figure 2.2.2

3 WDE and an additive Gaussian noise

3.1. Integral representations for WDEs. Following [8], [4], the WLSI (1.6) can be re-written (under certain conditions on f_{X_2}, f_{X_2} and ϕ) in terms of integral representations of the entropies $h^{\rm w}_{\phi}(X)$, $h^{\rm w}_{\phi_{\rm C}}(Y_1)$ and $h^{\rm w}_{\phi_{\rm S}}(Y_2)$: cf. Eqns (3.4), (3.5) below. Despite its cumbersome appearance, formulas (3.4) and (3.5) have an advantage: they does not include logarithms. (However, note condition (3.2).) Throughout the presentation in this section, the reader can notice persistent similarities with [4].

In this section we work with a two-variable WF $(x,y)\in\mathbb{R}\times\mathbb{R}\mapsto\rho(x,y)\geq 0$ and a number of reduced WDEs involving various integrals of ρ. Let Z and N be two independent RVs, where $N\sim\mathrm{N}(0,1)$ with standard normal PDF f^{No}, while Z has a PDF f_Z. Following [8] and [4], RV Z will represent a signal and N an (additive) Gaussian noise; RV Z will be a pre-cursor for $X=X_1+X_2$, Y_1 and Y_2. Given $\gamma>0$, $y\in\mathbb{R}$, set:

$$\begin{array}{l}\displaystyle \xi_Z(y,\gamma)=\frac{\displaystyle\int(y-t\sqrt{\gamma})yf^{\mathrm{No}}(y-t\sqrt{\gamma})f_Z(t)\mathrm{d}t}{\displaystyle\int f^{\mathrm{No}}(y-z\sqrt{\gamma})f_Z(z)\mathrm{d}z},\\ \displaystyle \eta_Z(y,\gamma)=\frac{\displaystyle\int(y-w\sqrt{\gamma})wf^{\mathrm{No}}(y-w\sqrt{\gamma})f_Z(w)\mathrm{d}w}{\displaystyle\int f^{\mathrm{No}}(y-z\sqrt{\gamma})f_Z(z)\mathrm{d}z},\\ \displaystyle \zeta_Z(x,y,\gamma)=\int_{-\infty}^{y}\rho(x,v)(v-x\sqrt{\gamma})x\frac{f_{Z,Z\sqrt{\gamma}+N}(x,v)}{f_{Z,Z\sqrt{\gamma}+N}(x,y)}\mathrm{d}v.\end{array}\tag{3.1}$$

Theorem 3. *Let X_1, X_2, N and N' be independent RVs, with $X=X_1+X_2$, where* (i) *X_j are with bounded and continuous PDFs f_{X_j} such that $\mathbb{E}|\log f_X(X)|<\infty$ and $\mathbb{E}|\log f_{X_j}(X_j)|<\infty$, $j=1,2$, and* (ii) *$N,N'\sim\mathrm{N}(0,1)$. Assume that for $Z=X_1$, $Z=X_2$ and $Z=X$, the conditional expectation $\mathbb{E}\left[f_Z\left(Z+\dfrac{N-N'}{\sqrt{\gamma}}\right)\Big|Z,N\right]$ is such that, for some integrable RV $\chi(Z,N)\geq 0$*

$$\left|\log\mathbb{E}\left[f_Z\left(Z+\frac{N-N'}{\sqrt{\gamma}}\right)\Big|Z,N\right]\right|\leq\chi(Z,N).\tag{3.2}$$

Next, consider a WF $(x,y)\in\mathbb{R}\times\mathbb{R}\mapsto\rho(x,y)$. Suppose that ρ is continuous and bounded, and $\forall\ x\in\mathbb{R}$, $\exists$ a limit $\phi(x)=\lim\limits_{y\to\pm\infty}\rho(x,y)$. Introduce additional WFs

$$\begin{array}{l}\rho_{\mathrm{C}}(x,y)=\rho(x\cos\vartheta,y),\ \rho_{\mathrm{S}}(x,y)=\rho(x\sin\vartheta,y),\ \text{with}\ \phi_{\mathrm{C/S}}(x)=\lim\limits_{y\to\pm\infty}\rho_{\mathrm{C/S}}(x,y),\\ \phi^*(v)=\int\rho(x,v)f_X(x)\mathrm{d}x,\ \phi_1^*(v)=\displaystyle\int\rho_{\mathrm{C}}(x,v)f_{Y_1}(x)\mathrm{d}x,\ \phi_2^*(v)=\int\rho_{\mathrm{S}}(x,v)f_{Y_2}(x)\mathrm{d}x\end{array}\tag{3.3}$$

where Y_2, Y_2 are as in (1.3).

Then, the WDE $h^{\mathrm{w}}_{\phi}(X)$ of the sum $X=X_1+X_2$ in the RHS of (1.6) *admits the representation*

$$\begin{array}{l}\displaystyle\int_0^\infty\frac{1}{2\sqrt{\gamma}}\,\mathbb{E}\Big[\zeta_X\Big(X,X\sqrt{\gamma}+N,\gamma\Big)\xi_X\Big(X\sqrt{\gamma}+N,\gamma\Big)\\ \qquad\qquad -\rho\Big(X,X\sqrt{\gamma}+N\Big)\eta_X\Big(X\sqrt{\gamma}+N,\gamma\Big)\Big]\mathrm{d}\gamma+h^{\mathrm{w}}_{\phi^*}(N).\end{array}\tag{3.4}$$

On the other hand, for $h^{\mathrm{w}}_{\phi_{\mathrm{C}}}(Y_1)$ and $h^{\mathrm{w}}_{\phi_{\mathrm{S}}}(Y_2)$ we have the respective formulas

$$\begin{array}{l}\displaystyle\int_0^\infty\frac{1}{2\sqrt{\gamma}}\,\mathbb{E}\Big[\zeta_{Y_j}\Big(Y_j,Y_j\sqrt{\gamma}+N,\gamma\Big)\xi_{Y_j}\Big(Y_j\sqrt{\gamma}+N,\gamma\Big)\\ \qquad\qquad -\rho\Big(Y_j,Y_j\sqrt{\gamma}+N\Big)\eta_{Y_j}\Big(Y_j\sqrt{\gamma}+N,\gamma\Big)\Big]\mathrm{d}\gamma+h^{\mathrm{w}}_{\phi_1^*}(N),\ j=1,2.\end{array}\tag{3.5}$$

The proof of Theorem 3 uses two technical assertions, Lemmas 3.1 and 3.2. They address the cases $\gamma = 0$ and $\gamma \to \infty$ (that is, the integration endpoints in (3.4) and (3.5)). For the definition of the weighted conditional and mutual entropies, see Eqns (1.11), (1.12) in [7].

Lemma 3.1. (Cf. Lemma 2.4 in [4].) *Let Z, U be independent RVs. Assume that U has a bounded and continous PDF $f_U \in \mathrm{C}^0(\mathbb{R}^d)$: $\int f_U(x)\mathrm{d}x = 1$ and $\operatorname{ess\,sup}[f_U(x), x \in \mathbb{R}^d] < +\infty$. The distribution of Z may have discrete and continous parts; we refer to the PMF $f_Z(x)$ relative to a reference measure $\nu(\mathrm{d}x)$. Next, suppose that a bounded WF $(x, y) \in \mathbb{R} \times \mathbb{R} \mapsto \rho(x, y)$ has been given and assume that $\mathbb{E}|\log f_Z(Z)| < +\infty$. Consider the weighted mutual entropy (WME) $i^{\mathrm{w}}_\rho(Z : \sqrt{\gamma}Z + U)$ between Z and $\sqrt{\gamma}Z + U$ where $\gamma > 0$ is a parameter. Then*

$$\lim_{\gamma \to 0} i^{\mathrm{w}}_\rho(Z : \sqrt{\gamma}Z + U) = 0. \tag{3.6}$$

Proof. According to the definition of the WME, for a pair of RVs Z, V with a conditional PDF $f_{V|Z}(y, x)$ we have an equality involving a weighted conditional entropy (WCE)

$$i^{\mathrm{w}}_\rho(Z : V) = h^{\mathrm{w}}_{\psi_Z}(Z) - h^{\mathrm{w}}_\rho(Z|V) \text{ where } \psi_Z(x) = \int \rho(x, y) f_{V|Z}(y|x)\mathrm{d}y.$$

Setting $V = \sqrt{\gamma}Z + U$, we can write a representation for the WCE:

$$\begin{aligned} h^{\mathrm{w}}_\rho(Z|\sqrt{\gamma}Z + U) = \int & \rho(x, y) f_U(y - \sqrt{\gamma}x) f_Z(x) \\ & \times \log\left[\frac{\int f_U(y - \sqrt{\gamma}z) f_Z(z) dz}{f_U(y - \sqrt{\gamma}x) f_Z(x)}\right] dy \nu(\mathrm{d}x). \end{aligned}$$

Using that ρ and f_U are bounded, with the help of the Lebesgue dominated convergence theorem we have that as $\gamma \to 0$, the ratio under the log converges to $[f_Z(x)]^{-1}$. Consequently,

$$\lim_{\gamma \to 0} h^{\mathrm{w}}_\rho(Z|\sqrt{\gamma}Z + U) = h^{\mathrm{w}}_{\psi^*_Z}(Z) \text{ where } \psi^*_Z(x) = \int \rho(x, y) f_U(y)\mathrm{d}y.$$

Moreover, with $\psi^*_{Z,\gamma}(x) = \displaystyle\int \rho(x, y) f_U(y - \sqrt{\gamma}x)\mathrm{d}y$, we introduce:

$$h^{\mathrm{w}}_{\psi^*_{Z,\gamma}}(Z) = -\int \rho(x, y) f_U(y - \sqrt{\gamma}x) f_Z(x) \log f_Z(x) \mathrm{d}y \nu(\mathrm{d}x).$$

At this stage we again apply the Lebesgue dominated convergence theorem and deduce that $\lim\limits_{\gamma \to 0} h^{\mathrm{w}}_{\psi^*_{Z,\gamma}}(Z) = h^{\mathrm{w}}_{\psi^*_Z}(Z)$. This leads to (3.6). $\square$

Lemma 3.2. (Cf. Lemma 4.1 in [4].) *Let Z, U and U' be independent RVs, Z with a PDF f_Z and U, U' with a PDF f_U. As before, consider a WF $(x, y) \in \mathbb{R} \times \mathbb{R} \mapsto \rho(x, y)$. Suppose that*

f_Z and ρ are continuous and bounded, and there exists a limit $\overline{\rho}(x)=\lim\limits_{y\to\infty}\rho(x,y)$. Next, assume that for some RV $\chi(Z,U)\geq 0$ with $\mathbb{E}\chi(Z,U)<\infty$, we have $\left|\log\mathbb{E}\left[f_Z\left(Z+\frac{U-U'}{\sqrt{\gamma}}\right)\Big|Z,U\right]\right|\leq\chi(Z,U)$. Then

$$\begin{aligned} &h^{\rm w}_{\overline{\rho}}(Z)=\lim_{\gamma\to\infty}\left[i^{\rm w}_{\rho}(Z:\sqrt{\gamma}Z+U)+h^{\rm w}_{\psi^*_{U,\gamma}}(U)\right]\ \textit{where}\\ &\quad h^{\rm w}_{\psi^*_{U,\gamma}}(U)=-\mathbb{E}\psi^*_{U,\gamma}(U)\log\, f_U(U)\ \text{and}\ \psi^*_{U,\gamma}(u)=\int\rho(x,u+\sqrt{\gamma}x)f_Z(x)dx.\end{aligned}\tag{3.7}$$

Proof. We can write

$$\begin{aligned}&i^{\rm w}_{\rho}(Z:\sqrt{\gamma}Z+U)+h^{\rm w}_{\psi^*_{U,\gamma}}(U)\\ &\quad=-\int\rho(x,u+\sqrt{\gamma}x)f_U(u)f_Z(x)\log\left[\int f_U(v)f_Z\left(x+\frac{u-v}{\sqrt{\gamma}}\right){\rm d}v\right]{\rm d}u{\rm d}x.\end{aligned}\tag{3.8}$$

Passing to the limit $\gamma\to\infty$, Eqn (3.8) yields (3.7), again owing to the Lebesgue dominated convergence theorem. $\Box$

Proof of Theorem 3. We again use Z as a substitute for RVs Y_1, Y_2 and $X=X_1+X_2$. Given $\gamma>0$, write the joint WDE for Z and $Z\sqrt{\gamma}+N$:

$$\begin{aligned}&h^{\rm w}_{\rho}(Z,Z\sqrt{\gamma}+N)=-\int\rho(x,x\sqrt{\gamma}+v)f^{\rm No}(v)f_Z(x)\log f^{\rm No}(v){\rm d}x{\rm d}v\\ &\quad-\int\rho(x,x\sqrt{\gamma}+v)f^{\rm No}(v)f_Z(x)\log f_Z(x){\rm d}x{\rm d}v=h^{\rm w}_{\psi^{(1)}_{Z,\gamma}}(Z)+h^{\rm w}_{\psi^{(2)}_{N,\gamma}}(N).\end{aligned}\tag{3.9}$$

Here and below, $\forall\ \gamma,\theta>0$,

$$\psi^{(1)}_{Z,\gamma}(x)=\int\rho(x,x\sqrt{\gamma}+v)f^{\rm No}(v){\rm d}v,\ \ \psi^{(2)}_{N,\theta}(v)=\int\rho(x,x\sqrt{\theta}+v)f_Z(x){\rm d}x,\tag{3.10}$$

with $h^{\rm w}_{\psi^{(2)}_{N,\theta}}(N)=-\int\left[\int\rho(x,x\sqrt{\theta}+v)f_Z(x){\rm d}x\right]f^{\rm No}(v)\log\, f^{\rm No}(v){\rm d}v$.

Moreover, according to Lemma 3.2 (with $U=N$, $U'=N'$), $\forall\ \epsilon>0$ we have

$$h^{\rm w}_{\overline{\rho}}(Z)=\int_{\epsilon}^{\infty}\frac{\rm d}{{\rm d}\gamma}\left[i^{\rm w}_{\rho}(Z:Z\sqrt{\gamma}+N)+h^{\rm w}_{\psi^{(2)}_{N,\gamma}}(N)\right]{\rm d}\gamma+i^{\rm w}_{\rho}(Z:Z\sqrt{\epsilon}+N)+h^{\rm w}_{\psi^{(2)}_{N,\epsilon}}(N).$$

To analyze the WDE $h^{\rm w}_{\psi^{(2)}_{N,\gamma}}(N)$, we use Lebesgue$'$s dominated convergence theorem. This yields

$$\lim_{\epsilon\to 0}h^{\rm w}_{\psi^{(2)}_{N,\epsilon}}(N)=h^{\rm w}_{\rho^*_N}(N)\ \ \text{where}\ \ \rho^*_N(v)=\int\rho(x,v)f_Z(x){\rm d}x.$$

In addition we get that the WME $i^{\rm w}_{\rho}(Z:Z\sqrt{\gamma}+N)$ is represented as the difference

$$h^{\rm w}_{\psi}(Z\sqrt{\gamma}+N)-h^{\rm w}_{\psi^{(2)}_{N,\gamma}}(N)\ \text{with}\ \psi(x)=\int\rho(x,v)f^{\rm No}(v-\sqrt{\gamma}x){\rm d}v.\tag{3.11}$$

Note that $\psi^{(2)}_{N,\gamma} = \psi^{*}_{U,\gamma}$; see (3.7). Furthermore, owing to Lemma 3.1 (with $U = N$), we write:

$$h^{\rm w}_{\overline{\rho}}(Z) = \int_0^{\infty} \frac{\mathrm{d}}{\mathrm{d}\gamma} h^{\rm w}_{\psi}(Z\sqrt{\gamma}+N)\mathrm{d}\gamma + h^{\rm w}_{\rho^*_N}(N), \text{ still with } \overline{\rho}(x) = \lim_{y\to\infty} \rho(x,y). \tag{3.12}$$

We are now going to analyze the derivative $\frac{\mathrm{d}}{\mathrm{d}\gamma} h^{\rm w}_{\psi}(Z\sqrt{\gamma}+N)$ representing it as

$$\begin{aligned}
&-\frac{\mathrm{d}}{\mathrm{d}\gamma}\int\int \rho(x,y) f^{\rm No}(y-x\sqrt{\gamma})\ f_Z(x)\log\left[\int f^{\rm No}(y-t\sqrt{\gamma})\ f_Z(t)\mathrm{d}t\right]\mathrm{d}x\mathrm{d}y \\
&= -\frac{1}{2\sqrt{\gamma}}\int\int \rho(x,y) f_Z(x) f^{\rm No}(y-x\sqrt{\gamma})(y-x\sqrt{\gamma})x \\
&\qquad \times\log\left[\int f^{\rm No}(y-t\sqrt{\gamma})\ f_Z(t)\mathrm{d}t\right]\mathrm{d}x\mathrm{d}y \\
&\quad -\frac{1}{2\sqrt{\gamma}}\int\int \rho(x,y) f_Z(x) f^{\rm No}(y-x\sqrt{\gamma}) \\
&\qquad \times\left[\frac{\int w(y-w\sqrt{\gamma}) f^{\rm No}(y-w\sqrt{\gamma}) f_Z(w)\mathrm{d}w}{\int f^{\rm No}(y-z\sqrt{\gamma}) f_Z(z)\mathrm{d}z}\right]\mathrm{d}x\mathrm{d}y.
\end{aligned} \tag{3.13}$$

The first integral in the RHS of (3.13) is done by parts. This leads to the following expression:

$$\begin{aligned}
&\frac{1}{2\sqrt{\gamma}}\int\int_{-\infty}^{y}\int \rho(x,v) f_Z(x) f^{\rm No}(v-x\sqrt{\gamma})(v-x\sqrt{\gamma})x\mathrm{d}x\mathrm{d}v \\
&\qquad\qquad \times\left[\frac{\int (y-t\sqrt{\gamma}) y f^{\rm No}(y-t\sqrt{\gamma}) f_Z(t)\mathrm{d}t}{\int f^{\rm No}(y-z\sqrt{\gamma}) f_Z(z)\mathrm{d}z}\right]\mathrm{d}y \\
&-\frac{1}{2\sqrt{\gamma}}\int\int \rho(x,y) f_Z(x) f^{\rm No}(y-x\sqrt{\gamma})\left[\frac{\int (y-w\sqrt{\gamma}) w f^{\rm No}(y-w\sqrt{\gamma}) f_Z(w)\mathrm{d}w}{\int f^{\rm No}(y-z\sqrt{\gamma}) f_Z(z)\mathrm{d}z}\right]\mathrm{d}x\mathrm{d}y.
\end{aligned}$$

Then, taking into account Eqn (3.1),

$$\begin{aligned}
\frac{\mathrm{d}}{\mathrm{d}\gamma} h^{\rm w}_{\psi}(Z\sqrt{\gamma}+N) &= \frac{1}{2\sqrt{\gamma}}\int\int \zeta(x,y,\gamma)\xi(y,\gamma) f_{Z,Z\sqrt{\gamma}+N}(x,y)\mathrm{d}x\mathrm{d}y \\
&\quad -\frac{1}{2\sqrt{\gamma}}\int\int \rho(x,y)\eta(y,\gamma) f_{Z,Z\sqrt{\gamma}+N}(x,y)\mathrm{d}x\mathrm{d}y \\
&= \frac{1}{2\sqrt{\gamma}}\ \mathbb{E}\Big\{\zeta_\rho(Z,Z\sqrt{\gamma}+N)\xi(Z\sqrt{\gamma}+N,\gamma) - \rho(Z,Z\sqrt{\gamma}+N)\eta(Z\sqrt{\gamma}+N,\gamma)\Big\}.
\end{aligned} \tag{3.14}$$

□

3.2. WLSI for a WF close to a constant. Concluding this section, we analyze the WLSI (1.6) when the WF ϕ lies in vicinity of a constant $\overline{\phi}$ (and hence, is bounded). Given independent RVs X_1, X_2 with PDFs f_{X_1} and f_{X_2}, we refer to Y_1 and Y_2 as $Y_1 = X_1/\cos\alpha$, $Y_2 = X_2/\sin\alpha$ where $\alpha \in [-\pi, \pi)$ is as in Eqn (1.3). For $Z = Y_1, Y_2$ or $X_1 + X_2 = Y_1\cos\alpha + Y_2\sin\alpha$, set:

$$M(Z;\gamma) = \mathbb{E}\Big[\big|Z - \mathbb{E}\big(Z\big|Z\sqrt{\gamma}+N\big)\big|^2\Big] \tag{3.15}$$

and suppose that for the above choices of RV Z:

$$\mathbb{E}|\ln f_Z(Z)| < \infty. \tag{3.16}$$

We also need to assume a uniform integrability condition (3.2): for an independent triple Z, N, N' where $N, N' \sim \mathrm{N}(0,1)$, there exists an integrable RV $\chi(Z,N) \geq 0$ such that

$$\left| \log \mathbb{E}\left[f_Z\left(Z + \frac{N - N'}{\sqrt{\gamma}} \right) \Big| Z, N \right] \right| \leq \chi(Z,N). \tag{3.17}$$

According to formula (4.5) in [4], for these choices of Z, the standard SDE $h(Z)$ is represented as

$$h(Z) = h(N) + \frac{1}{2} \int \left[M(Z,\gamma) - \mathbf{1}(\gamma > 1)\frac{1}{\gamma} \right] \mathrm{d}\gamma. \tag{3.18}$$

Furthermore, as follows from the proof of Theorem 4.1 in [4] (see [4], Eqn (4.8)), for any $\widetilde{\alpha} \in [-\pi, \pi]$ (including $\widetilde{\alpha} = \alpha$, the value from (1.3)),

$$M(Y_1 \cos\widetilde{\alpha} + Y_2 \sin\widetilde{\alpha}, \gamma) \geq M(Y_1,\gamma)(\cos\widetilde{\alpha})^2 + M(Y_2,\gamma)(\sin\widetilde{\alpha})^2.$$

For $\widetilde{\alpha} = \alpha$, this becomes

$$M(X_1 + X_2, \gamma) \geq M(Y_1,\gamma)(\cos\alpha)^2 + M(Y_2,\gamma)(\sin\alpha)^2. \tag{3.19}$$

Now we are in position to establish Theorem 4 below. As before, we refer to $Z = Y_1, Y_2$ or $X_1 + X_2 = Y_1\cos\alpha + Y_2 \sin\alpha$.

Theorem 4. *Let $\gamma_0 > 0$ be is a point of continuity of $M(Z,\gamma)$, $Z = Y_1, Y_2, X_1 + X_2$. Suppose that there exists $\delta > 0$ such that*

$$M(X_1 + X_2, \gamma_0) \geq M(Y_1,\gamma_0)(\cos\alpha)^2 + M(Y_2,\gamma_0)(\sin\alpha)^2 + \delta. \tag{3.20}$$

Also assume (3.16) *and* (3.17).

Then there exists $\epsilon = \epsilon(\gamma_0, \delta, f_{X_1}, f_{X_2})$ with the following property. Let function $x \in \mathbb{R} \mapsto \phi(x) \geq 0$ be such that $\left|\phi(x) - \overline{\phi}\right| \leq \epsilon$, $\forall\ x$, for a constant $\overline{\phi} > 0$. Then the WLSI (1.6) *with the WF ϕ holds true.*

Proof of Theorem 4. According to Theorem 1, to prove the WSLI (1.6), we only need to check that

$$(\cos\alpha)^2 h^{\mathrm{w}}_{\phi_{\mathrm{C}}}(Y_1) + (\sin\alpha)^2 h^{\mathrm{w}}_{\phi_{\mathrm{S}}}(Y_2) \leq h^{\mathrm{w}}_{\phi}(Y_1\cos\alpha + Y_2\sin\alpha).$$

For a constant WF $\overline{\phi}$, the following inequality is valid (see Ref [4], Lemma 4.2 or Ref [8], Eqns (9) and (10))

$$(\cos\alpha)^2 h^{\mathrm{w}}_{\overline{\phi}}(Y_1) + (\sin\alpha)^2 h^{\mathrm{w}}_{\overline{\phi}}(Y_2) \leq h^{\mathrm{w}}_{\overline{\phi}}(Y_1\cos\alpha + Y_2\sin\alpha).$$

Next, Theorem 4.1 from [4] (applicable because of (3.16) and (3.17)) implies that under condition (3.20), for ϵ small enough

$$(\cos\alpha)^2 h^{\rm w}_{\overline{\phi}}(Y_1) + (\sin\alpha)^2 h^{\rm w}_{\overline{\phi}}(Y_2) + \epsilon \leq h^{\rm w}_{\overline{\phi}}(Y_1\cos\alpha + Y_2\sin\alpha).$$

Define $\varphi(x) = |\phi(x) - \overline{\phi}|$. It remains to check that

$$h^{\rm w}_{\varphi}(Y_1) \leq \epsilon/3, h^{\rm w}_{\varphi}(Y_2) \leq \epsilon/3, h^{\rm w}_{\varphi}(Y_1\cos\alpha + Y_2\sin\alpha) \leq \epsilon/3.$$

But this inequality immediately follows, owing to (3.16). This competes the proof of Theorem 4. $\square$

The statement of Theorem 4 can be made more efficient for given PDFs f_{X_1} and f_{X_1}. As an example, consider the case where RVs X_1, X_2 are normal and WF $\phi \in C^2$.

Lemma 3.3. *Let RVs* $X_i \sim$*N*$(\mu_i, \sigma_i^2), i = 1, 2$ *be independent, and* $X = X_1 + X_2 \sim$ N$(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$. *Suppose that WF* $x \in \mathbb{R} \mapsto \phi(x) \geq 0$ *is twice contiuously differentiable and slowly varying in the sense that* $\forall$ x,

$$|\phi''(x)| \leq \epsilon\phi(x), \ \ |\phi(x) - \overline{\phi}| < \epsilon, \tag{3.21}$$

where $\epsilon > 0$ *and* $\overline{\phi} > 0$ *are constants. Then there exists* $\epsilon_0 = \epsilon_0(\mu_0, \mu_1, \sigma_0^2, \sigma_1^2) > 0$ *such that for any* $0 < \epsilon \leq \epsilon_0$, *the WLSI* (2.3) *with the WF* ϕ *holds true.*

Proof of Lemma 3.3. Let α be as in Eqn (1.3); to check (2.3), we use Stein's formula: for $Z \sim$ N(μ, σ^2),

$$\mathbb{E}\Big[Z^2\phi(Z)\Big] = \sigma^2\mathbb{E}\Big[\phi(Z)\Big] + \sigma^4\mathbb{E}\Big[\phi''(Z)\Big]. \tag{3.22}$$

Owing to the inequality $|\phi(x) - \overline{\phi}| < \epsilon$,

$$\alpha < \alpha_0 = \tan^{-1}\Big(\exp\Big\{(\overline{\phi} + \epsilon)^2[h_+(X_2) - h_-(X_1)] - (\overline{\phi} - \epsilon)^2[h_+(X_1) - h_-(X_2)]\Big\}\Big).$$

Here

$$h_\pm(X_i) = -\mathbb{E}\Big[\mathbf{1}(X_i \in A^i_\pm)\log f^{\rm No}_{X_i}(X_i)\Big], \ \ i = 1, 2,$$

and

$$A^i_+ = \Big\{x \in \mathbb{R} : \ f^{\rm No}_i(x) < 1\Big\}, \ \ A^i_- = \Big\{x \in \mathbb{R} : \ f^{\rm No}_i(x) > 1\Big\}.$$

Evidently, under conditions $|\phi'(x)|, |\phi''(x)| \leq \epsilon\phi(x)$ we have that $\alpha_0 < \frac{\pi}{2} - \epsilon$ and $0 < \epsilon < (\sin\alpha)^2, (\cos\alpha)^2 < 1 - \epsilon < 1$. We claim that inequality (2.3) is satisfied with ϕ replaced by $\overline{\phi}$

and added $\delta > 0$:

$$\begin{aligned}&\Big[\log\Big(2\pi(\sigma_1^2+\sigma_2^2)\Big)\Big]\overline{\phi}+\frac{\log e}{\sigma_1^2+\sigma_2^2}\,\overline{\phi}\mathbb{E}\big[X^2\big]\\&\geq(\cos\alpha)^2\left[\log\frac{2\pi\sigma_1^2}{(\cos\alpha)^2}\right]\overline{\phi}+\frac{(\cos\alpha)^2\log e}{\sigma_1^2}\,\overline{\phi}\mathbb{E}[X_1^2]\\&\quad+(\sin\alpha)^2\left[\log\frac{2\pi\sigma_2^2}{(\sin\alpha)^2}\right]\overline{\phi}+\frac{(\sin\alpha)^2\log e}{\sigma_2^2}\,\overline{\phi}\mathbb{E}[X_2^2]+\delta.\end{aligned}\tag{3.23}$$

Here $\delta > 0$ is calculated through ϵ and increases to a limit $\delta_0 > 0$ as $\epsilon \to 0$.

Indeed, strict concavity of $\log\, y$ for $y > 0$ implies that

$$\Big[\log\Big(2\pi(\sigma_1^2+\sigma_2^2)\Big)\Big]\geq(\cos\alpha)^2\left[\log\frac{2\pi\sigma_1^2}{(\cos\alpha)^2}\right]\overline{\phi}+(\sin\alpha)^2\left[\log\frac{2\pi\sigma_2^2}{(\sin\alpha)^2}\right]+\delta.$$

On the other hand,

$$\frac{1}{\sigma_1^2+\sigma_2^2}\,\overline{\phi}\mathbb{E}\big[X^2\big]=\frac{(\cos\alpha)^2}{\sigma_1^2}\,\overline{\phi}\mathbb{E}[X_1^2]+\frac{(\sin\alpha)^2}{\sigma_2^2}\,\overline{\phi}\mathbb{E}[X_2^2].$$

Now, if we look at Eqn (2.3) with WF ϕ then, owing to Eqn (3.23) it suffices to verify that

$$\begin{aligned}&\Big[\log\Big(2\pi(\sigma_1^2+\sigma_2^2)\Big)\Big]\mathbb{E}(\phi(X)-\overline{\phi})+\frac{\log e}{\sigma_1^2+\sigma_2^2}\,\mathbb{E}\big[X^2(\phi(X)-\overline{\phi})\big]\\&-(\cos\alpha)^2\left[\log\frac{2\pi\sigma_1^2}{(\cos\alpha)^2}\right]\mathbb{E}(\phi(X_1)-\overline{\phi})+\frac{(\cos\alpha)^2\log e}{\sigma_1^2}\,\mathbb{E}[X_1^2(\phi(X_1)-\overline{\phi})]\\&-(\sin\alpha)^2\left[\log\frac{2\pi\sigma_2^2}{(\sin\alpha)^2}\right]\mathbb{E}(\phi(X_2)-\overline{\phi})+\frac{(\sin\alpha)^2\log e}{\sigma_2^2}\,\mathbb{E}[X_2^2(\phi(X_2)-\overline{\phi})]<\delta.\end{aligned}\tag{3.24}$$

We check this by a brute force, claiming that each term in (3.24) has the absolute value $< \delta/6$ when ϵ is small enough. For the terms containing $\mathbb{E}(\phi(Z)-\overline{\phi})$, $Z = X, X_1, X_2$, this follows since $|\phi(x)-\overline{\phi}| < \epsilon$.

The terms containing factor $\mathbb{E}\big[Z^2(\phi(Z)-\overline{\phi})\big]$ we use Stein‘s formula (3.22) and the condition that $|\phi''(x)| \leq \epsilon\phi(x)$. $\square$

Similar assertions can be established for other examples of PDFs f_{X_1} and f_{X_2}.

Acknowledgements – YS thanks the Math Department, Penn State University, for the financial support and hospitality during the academic year 2014-5. SYS thanks the CAPES PNPD-UFSCAR Foundation for the financial support in the year 2014-5. SYS thanks the Federal University of Sao Carlos, Department of Statistics, for hospitality during the year 2014-5. MK acknowledge the support of thew National Research University Higher School of Economics in the framework of implementation of the 5-100 Pregramme Roadmap.

References

[1] M. Belis and S. Guiasu. A Quantitative and qualitative measure of information in cybernetic systems. *IEEE Trans. on Inf. Theory*, **14** (1968), 593–594.

[2] A. Clim. Weighted entropy with application. *Analele Universităţii Bucureşti, Matematică*, Anul **LVII** (2008), 223-231.

[3] T. Cover and J. Thomas. *Elements of Information Theory.* New York: Wiley, 2006.

[4] M. Kelbert and Y. Suhov. Continuity of mutual entropy in the limiting signal-to-noise ratio regimes. In: *Stochastic Analysis*, Springer-Verlag: Berlin (2010), 281–299.

[5] M. Kelbert and Y. Suhov. *Information Theory and Coding by Example.* Cambridge: Cambridge University Press, 2013.

[6] E.H. Lieb. Proof of entropy conjecture of Wehrl. *Commun. Math. Phys.*, **62** (1978), 35–41.

[7] Y. Suhov and S. Yasaei Sekeh. Simple inequalities for weighted entropies. arXiv 1409.4102.

[8] S. Verdú and D. Guo. A simple proof of the entropy -power inequality. *IEEE Transaction on Information Theory*, **52**, No. 5 (2006), 2165–2166.